arXiv:math/0305074v3 [math.NT] 1 Dec 2003

ON THE CAUCHY PROBLEM FOR DIFFERENTIAL EQUATIONS IN A BANACH SPACE OVER THE FIELD OF p-ADIC NUMBERS. I. [1]

MYROSLAV L. GORBACHUK and VALENTYNA I. GORBACHUK

ABSTRACT. For the Cauchy problem for an operator differential equation of the form $y'(z) = Ay(z)$, where A is a closed linear operator on a Banach space over the field of p-adic numbers, the criterion of well-posedness in the class of locally analytic vector-functions is established. It is shown how the Cauchy-Kovalevskaya theorem for p-adic partial differential equations may be obtained as a particular case from this criterion.

1. Let $\mathfrak{B}$ be a Banach space with norm $\|\cdot\|$ over the completion $\Omega = \Omega_p$ of an algebraic closure of the field of p-adic numbers [1 - 3] (p is prime), and let A be a closed linear operator on $\mathfrak{B}$, that is, the convergences $\mathcal{D}(A) \ni x_n \to x$ and $Ax_n \to y$ $(n \to \infty)$ in $\mathfrak{B}$ imply the inclusion $x \in \mathcal{D}(A)$ and the equality $Ax = y$ ($\mathcal{D}(\cdot)$ is the domain of an operator).

For a number $\alpha > 0$, we put

$$E_\alpha(A) = \Big\{x \in \bigcap_{n\in\mathbb{N}_0=\{0,1,2,\dots\}} \mathcal{D}(A^n) \Big| \exists c = c(x) > 0 \quad \forall k \in \mathbb{N}_0 \quad \|A^k x\| \le c\alpha^k\Big\}.$$

The linear space $E_\alpha(A)$ is a Banach space with respect to the norm

$$\|x\|_\alpha = \sup_{n\in\mathbb{N}_0} \frac{\|A^n x\|}{\alpha^n}.$$

The set

$$E(A) = \bigcup_{\alpha>0} E_\alpha(A)$$

is endowed with the inductive limit topology of the Banach spaces $E_\alpha(A)$:

$$E(A) = \operatorname*{ind\,lim}_{\alpha\to\infty} E_\alpha(A).$$

By the closed graph theorem, $E(A)$ coincides with $\mathfrak{B}$ if and only if $\mathcal{D}(A) = \mathfrak{B}$. The elements of $E(A)$ are called entire vectors of exponential type for the operator A. Define the type $\sigma(x; A)$ of a vector $x \in E(A)$ as

$$\sigma(x; A) = \inf\{\alpha > 0 : x \in E_\alpha(A)\} = \varlimsup_{n\to\infty} \|A^n x\|^{\frac{1}{n}}.$$

Thus, the equality $\sigma(x; A) = \sigma$ means that for an arbitrary $\varepsilon > 0$, there exists a constant $c_\varepsilon = c_\varepsilon(x) > 0$ such that

$$\forall n \in \mathbb{N}_0 \quad \|A^n x\| \le c_\varepsilon(\sigma + \varepsilon)^n,$$

---

[1] *Mathematics Subject Classification.* Primary 34G10

Key words and phrases. Field of p-adic numbers, differential equation in a Banacc space, entire vector of exponential type, Cauchy problem, Cauchy-Kovalevskaya theorem.

Supported by CRDF (Project UM 1-2421-KV-02)

and

$$\lim_{i\to\infty} \frac{\|A^{n_i}x\|}{(\sigma-\varepsilon)^{n_i}} = \infty$$

for some subsequence $n_i \to \infty$ when $i \to \infty$. In the case, where the operator A is bounded, the type of a vector $x \in \mathfrak{B} = E(A)$ does not exceed the norm of A:

$$\forall x \in \mathfrak{B} \quad \sigma(x; A) \le \|A\|.$$

2. The object of consideration now is a power series

$$y(z) = \sum_{n=0}^{\infty} c_n z^n, \quad c_n \in \mathfrak{B}, \quad z \in \Omega. \tag{1}$$

For such a series the convergence radius is determined by the formula

$$r = r(y) = \frac{1}{\overline{\lim}_{n\to\infty} \sqrt[n]{\|c_n\|}}. \tag{2}$$

In the open disk $d(r^-; \Omega) = \{z \in \Omega : |z|_p < r\}$, the series (1) defines a vector-function $y(z)$ with values in $\mathfrak{B}$ ($|\cdot|_p$ is the p-adic valuation on Ω).

Denote by $\mathfrak{A}_{loc}(\mathfrak{B})$ the set of all vector-functions $y(z)$ which are represented by a series of the form (1) with $r(y) > 0$. It is obvious that $\mathfrak{A}_{loc}(\mathfrak{B})$ is a vector space over Ω. We call its elements locally analytic vector-functions. The convergence $y_n \to y$ $(n \to \infty)$ in $\mathfrak{A}_{loc}(\mathfrak{B})$ means that there exists a number $\delta > 0$ such that $r(y_n) \ge \delta$, for any $n \in \mathbb{N}$, and for an arbitrary $\varepsilon \in (0, \delta)$,

$$\sup_{|z|_p \le \delta - \varepsilon} \|y_n(z) - y(z)\| \to 0, \ n \to \infty.$$

Let $y \in \mathfrak{A}_{loc}(\mathfrak{B})$. Its derivatives are defined as

$$y^{(k)}(z) = \sum_{n=0}^{\infty} (n+1)\dots(n+k) c_{n+k} z^n, \quad k \in \mathbb{N}.$$

It follows from (2) that

$$r(y^{(k)}) \ge r(y).$$

It is easily checked also that if $z \to 0$, then

$$y(z) \to y(0) = c_0, \text{ and } \frac{y^{(k)}(z) - y^{(k)}(0)}{z} \to y^{(k+1)}(0) = c_{k+1}(k+1)! \tag{3}$$

in the topology of $\mathfrak{B}$.

3. Let us consider the Cauchy problem

$$\left\{ \begin{array}{rcl} \dfrac{dy(z)}{dz} & = & Ay(z) \\ y(0) & = & y_0, \end{array} \right. \tag{4}$$

where A is a closed linear operator on $\mathfrak{B}$. We say that a vector-function $y(z)$ from $\mathfrak{A}_{loc}(\mathfrak{B})$ is a solution of problem (4) if $y(z) \in \mathcal{D}(A)$ for $z \in d(r(y)^-; \Omega)$ and satisfies (4) in this disk.

Theorem 1. *In order that problem (4) have a solution in $\mathfrak{A}_{loc}(\mathfrak{B})$, it is necesary and sufficient that $y_0 \in E(A)$; moreover $\sigma(y_0; A)r(y) = p^{-\frac{1}{p-1}}$. The problem (4) is well-posed, that is, the solution is unique, anf if a sequence of initial data $y_{n,0} \in E(A)$ converges to y_0 in $E(A)$, then the sequence of the corresponding solutions $y_n(z) \in \mathfrak{A}_{loc}(\mathfrak{B})$ converges to $y(z)$ in the space $\mathfrak{A}_{loc}(\mathfrak{B})$.*

Proof. Suppose that

$$y(z) = \sum_{k=0}^{\infty} c_k z^k, \quad c_k \in \mathfrak{B}, \quad z \in d(r(y)^-; \Omega),$$

is a solution of (4). Then $c_k \in \mathcal{D}(A),\ k \in \mathbb{N}_0$. Indeed, $c_0 = y(0) = y_0 \in \mathcal{D}(A)$. Since

$$\mathcal{D}(A) \ni \frac{y(z) - c_0}{z} = \sum_{k=0}^{\infty} c_k z^{k-1} \to c_1,$$

and

$$\frac{A(y(z) - y_0)}{z} = \frac{y'(z) - y'(0)}{z} \to 2c_2$$

as $z \to 0$, we have because of closure of the operator A that $c_1 \in \mathcal{D}(A)$, and $Ac_1 = 2c_2$. Using (3), we get by induction that

$$\forall k \in \mathbb{N} \quad c_k = \frac{Ac_{k-1}}{k} \in \mathcal{D}(A),$$

hence,

$$\forall k \in \mathbb{N}\ y_0 \in \mathcal{D}(A^k),\ \text{and}\ A^k y_0 = k! c_k.$$

In view of (2),

$$\frac{1}{r(y)} = \overline{\lim_{n\to\infty}} \sqrt[n]{\frac{\|A^n y_0\|}{|n!|_p}}.$$

Taking into account the equality

$$\lim_{n\to\infty} \sqrt[n]{|n!|_p} = p^{-\frac{1}{p-1}}$$

(see [1]), we obtain

$$\overline{\lim_{n\to\infty}} \sqrt[n]{\|A^n y_0\|} = \frac{p^{-\frac{1}{p-1}}}{r(y)},$$

whence

$$\forall n \in \mathbb{N}_0 \quad \|A^n y_0\| \le c\alpha^n,$$

where $0 < c = \text{const},\ \alpha = \frac{p^{-\frac{1}{p-1}}}{r(y)}$. So, $y_0 \in E(A)$, and $\sigma(y_0; A)r(y) = p^{-\frac{1}{p-1}}$.

Conversely, let y_0 be an entire vector of exponential type for the operator A with $\sigma(y_0; A) = \sigma$. Then the series

$$y(z) = \sum_{k=0}^{\infty} \frac{A^k y_0}{k!} z^k \tag{5}$$

is convergent in the disk $d(r^-; \Omega)$, where

$$r = r(y) = \left(\overline{\lim_{n\to\infty}} \sqrt[n]{\frac{\|A^n y_0\|}{|n!|_p}}\right)^{-1} = \frac{\lim_{n\to\infty} \sqrt[n]{|n!|_p}}{\overline{\lim}_{n\to\infty} \sqrt[n]{\|A^n y_0\|}} = \frac{p^{-\frac{1}{p-1}}}{\sigma}.$$

We shall prove now that if $z \in d(r^-;\Omega)$, then $y(z) \in \mathcal{D}(A)$. Really, since every component of series (5) belongs to $\mathcal{D}(A)$, the sums $S_n(z) = \sum\limits_{k=0}^{n} \dfrac{A^k y_0}{k!} z^k$ belong to $\mathcal{D}(A)$, too. For $z \in d(r^-;\Omega)$, the sequence $S_n(z)$ converges to $y(z)$ $(n \to \infty)$ in the topology of $\mathfrak{B}$. As $\sigma(Ay_0; A) = \sigma(y_0; A) = \sigma$, the sequence $AS_n(z) = \sum\limits_{k=0}^{n} \dfrac{A^{k+1} y_0}{k!} z^k$, $n \in \mathbb{N}$, converges in $\mathfrak{B}$ $(n \to \infty)$ in the same disk $d(r^-;\Omega)$. Since the operator A is closed, we have that $y(z) \in \mathcal{D}(A)$ when $z \in d(r^-;\Omega)$.

The formal differentiation of series (5) verifies that $y(z)$ satisfies (4). Thus, the vector-function (5) is a solution of problem (4).

It remains to check the well-posedness of problem (4). Assume that $y_{n,0} \to y_0$ $(n \to \infty)$ in $E(A)$. This means that there exists a number $\alpha > 0$ such that $y_{n,0} \in E_\alpha(A)$ for sufficiently large n, and $\|y_{n,0} - y_0\|_\alpha \to 0$ as $n \to \infty$. It follows from the above proof of sufficiency that

$$r(y_n) \geq \frac{p^{-\frac{1}{p-1}}}{\alpha}.$$

So, we may take $\delta = \dfrac{p^{-\frac{1}{p-1}}}{\alpha}$, and to complete the proof, we need only show that for an arbitrary fixed $\varepsilon \in (0,1)$, $\|y_n(z) - y(z)\| \to 0$ $(n \to \infty)$ uniformly in the disk $d((1-\varepsilon)\delta^-;\Omega)$. We have

$$\|y_n(z) - y(z)\| = \left\| \sum_{k=0}^{\infty} \frac{A^k(y_{n,0} - y_0) z^k}{k!} \right\| \leq \sum_{k=0}^{\infty} \frac{\|A^k(y_{n,0} - y_0)\| |z|_p^k}{|k!|_p} =$$

$$\sum_{k=0}^{\infty} \frac{\|A^k(y_{n,0} - y_0)\|}{\alpha^k} \cdot \frac{\alpha^k |z|_p^k}{|k!|_p} \leq \sup_k \frac{\|A^k(y_{n,0} - y_0)\|}{\alpha^k} \cdot \sum_{k=0}^{\infty} \frac{\alpha^k |z|_p^k}{|k!|_p} \leq \|y_{n,0} - y_0\|_\alpha \sum_{k=0}^{\infty} \alpha^k (1-\varepsilon)^k \frac{p^{-\frac{k}{p-1}}}{\alpha^k |k!|_p}.$$

Taking into account that

$$\frac{1}{|k!|_p} \leq p^{\frac{k}{p-1}}$$

(see [2]), we arrive at the inequality

$$\|y_n(z) - y(z)\| \leq \varepsilon^{-1} \|y_{n,0} - y_0\|_\alpha,$$

and therefore problem (4) is well-posed. $\square$

It is seen from the proof of Theorem 1 that the series $\sum\limits_{k=0}^{\infty} \dfrac{A^k y_0}{k!} z^k$ is convergent for all $z \in \Omega$ if and only if $y_0 \in \bigcap\limits_{\alpha > 0} E_\alpha(A)$.

Corollary 1. *If the operator A is bounded, then for any $y_0 \in \mathfrak{B}$, the Cauchy problem (4) is well-posed in the class $\mathfrak{A}_{loc}(\mathfrak{B})$.*

Remark 1. *If $\mathfrak{B}$ is a Banach space over the field $\mathbb{C}$ of complex numbers, then, as was shown in [4], problem (4) is well-posed in the class of locally analytic vector-functions if and only if y_0 is an analytic vector for the operator A, that is, $y_0 \in \bigcap\limits_{n \in \mathbb{N}_0} \mathcal{D}(A^n)$, and*

$$\exists \alpha > 0 \quad \exists c > 0 \quad \forall k \in \mathbb{N}_0 \quad \|A^k y_0\| \leq c \alpha^k k!.$$

In this case, in order that problem (4) be well-posed in the class of entire vector-functions of exponential type, it is necessary and sufficient that $y_0 \in E(A)$.

4. In this section we show how the existence and uniqueness theorem for the Cauchy problem for partial differential equations over a non-archimedean field of characteristic zero (see [5]) may be obtained from the above result.

Let $\mathcal{A}_\rho$ be the space of Ω-valued functions $f(x)$ analytic on the n-dimensional disk

$$d(\rho^+;\Omega^n) = \left\{ x = (x_1, \dots x_n) \in \Omega^n : |x|_p = \left(\sum_{i=1}^n |x_i|_p^2 \right)^{1/2} \le \rho \right\}.$$

This means that

$$f(x) = \sum_\alpha f_\alpha x^\alpha, \quad f_\alpha \in \Omega, \quad \lim_{|\alpha|\to\infty} |f_\alpha|_p \rho^{|\alpha|} = 0,$$

where $\alpha = (\alpha_1, \dots \alpha_n),\ \alpha_i \in \mathbb{N}_0,\ |\alpha| = \alpha_1 + \dots + \alpha_n$.

The space $\mathcal{A}_\rho$ is a non-archimedean Banach space with respect to the norm

$$\|f\|_\rho = \sup_\alpha |f_\alpha|_p \rho^{|\alpha|}.$$

It is clear that the differential operators

$$\frac{\partial f}{\partial x_j} = \sum_\alpha \alpha_j f_\alpha x_1^{\alpha_1} \dots x_j^{\alpha_j - 1} \dots x_n^{\alpha_n},\ j = 1, \dots, n,$$

are bounded in $\mathcal{A}_\rho$, and

$$\left\| \frac{\partial f}{\partial x_j} \right\|_\rho = \max_\alpha |f_\alpha \alpha_j|_p \rho^{|\alpha|-1} \le \frac{1}{\rho} \max_\alpha |f_\alpha|_p \rho^{|\alpha|} = \frac{1}{\rho} \|f\|_\rho.$$

The multiplication map

$$G : f \mapsto fg,$$

where $g \in \mathcal{A}_\rho$, is bounded in $\mathcal{A}_\rho$, too, and

$$\|G\| = \|g\|_\rho.$$

Indeed, let $f(x) = \sum\limits_\alpha f_\alpha x^\alpha,\ g(x) = \sum\limits_\alpha g_\alpha x^\alpha,\ \alpha = (\alpha_1, \dots, \alpha_n)$. Then

$$f(x)g(x) = \sum_\alpha c_\alpha x^\alpha,$$

where

$$c_\alpha = \sum_{0 \le i \le \alpha} f_i g_{\alpha - i} = \sum_{i_1=0}^{\alpha_1} \cdots \sum_{i_n=0}^{\alpha_n} f_{i_1,\dots,i_n} g_{\alpha_1 - i_1, \dots, \alpha_n - i_n} \quad (i = (i_1, \dots, i_n)).$$

So,

$$\|fg\|_\rho = \sup_\alpha \max_{0 \le i \le \alpha} |f_i|_p |g_{\alpha-i}|_p \rho^{|i|} \rho^{|\alpha| - |i|} \le \|f\|_\rho \|g\|_\rho.$$

We pass now to the Cauchy problem

$$\left\{ \begin{array}{rcl} \dfrac{\partial u(t,x)}{\partial t} & = & \sum\limits_{|\beta|=0}^{n} a_\beta(x) D^\beta u(t,x) \\ u(0,x) & = & \varphi(x), \end{array} \right. \tag{6}$$

where $a_\beta(x) \in \mathcal{A}_\rho,\ \varphi(x) \in \mathcal{A}_\rho,\ D^\beta = \dfrac{\partial^{|\beta|}}{\partial x_1^{\beta_1} \dots \partial x_n^{\beta_n}}$.

In the space $\mathcal{A}_\rho$ we define the operator A as follows:

$$f \mapsto Af = \sum_{|\beta|=0}^{n} a_\beta D^\beta f.$$

The relations

$$\Big\| \sum_{|\beta|=0}^{n} a_\beta D^\beta f \Big\|_\rho \leq \max_\beta \|a_\beta D^\beta f\|_\rho \leq \max_\beta \|a_\beta\|_\rho \|D^\beta f\|_\rho \leq \max_\beta \{\rho^{-|\beta|} \|a_\beta\|_\rho\} \|f\|_\rho$$

show that the operator A is bounded, and

$$\|A\| \leq \max_\beta \{\rho^{-|\beta|} \|a_\beta\|_\rho\}.$$

It follows from Corollary 1 that problem (6) is well-posed in $\mathfrak{A}_{loc}(\mathcal{A}_\rho)$ in the disk

$$\left\{ t \in \Omega : |t|_p < \frac{p^{-\frac{1}{p-1}}}{\max\limits_\beta \{\rho^{-|\beta|} \|a_\beta\|_\rho\}} \right\}.$$

It should be noted that Theorem 1 is valid also in the case where $\mathfrak{B}$ is a Banach space over an arbitrary non-archimedean field K of characteristic zero. But then the role of $p^{-\frac{1}{p-1}}$ is played by a certain constant $b = b(K)$ such that $\dfrac{1}{|n!|_K} \leq b^n$ (see, for instance, [6]).

References

[1] Koblitz N. *p-adic numbers, p-adic Analysis, and Zeta-functions*, Springer-Verlag, New York-Heidelberg-Berlin, 1977; Russian edition: Mir, Moscow, 1982.

[2] Dwork B., Gerotto G., and Sullivan F.J. *An Introduction to G-functions*, Princeton University Press, Princeton-New Jersey, 1994.

[3] Schikhof W.H. *Ultrametric Calculus, An Introdunction to p-adic Analysis*, Cambridge University Press, London-New York-New Rochelle-Melburnr-Sydney, 1984.

[4] Gorbachuk M. L. *On analytic solutions of differential-operator equations*, Ukrain. Mat. Zh. **52** (2000), no. 2, 596-607.

[5] Khrennikov A.Yu. *Mathematical methods of non-archimedean physics*, Uspehi. Mat. Nauk. **45** (1990), no. 4, 79-110.

[6] Borevich Z.I., Shafarevich I.R. *Number Theory*, Nauka, Moscow, 1985.

Institute of Mathematics
National Academy of Sciences of Ukraine
3 Tereshchenkivs'ka
Kyiv 01601, Ukraine
E-mail: imath@horbach.kiev.ua